

## Role of regulatory miRNAs of the Wnt/ $\beta$ -catenin signaling pathway in tumorigenesis of breast cancer

Article (Accepted Version)

Rahmani, Farzad, Tadayyon Tabrizi, Ayda, Hashemian, Pedram, Alijannejad, Sajede, Rahdar, Hossein Ali, Ferns, Gordon A, Hassanian, Seyed Mahdi, Shahidsales, Soodabeh and Avan, Amir (2020) Role of regulatory miRNAs of the Wnt/  $\beta$ -catenin signaling pathway in tumorigenesis of breast cancer. *Gene*, 754. a144892. ISSN 0378-1119

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/93322/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

### **Copyright and reuse:**

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

# **Role of regulatory miRNAs of the Wnt/ $\beta$ -catenin signaling pathway in tumorigenesis of breast cancer**

Farzad Rahmani<sup>1,2</sup>, Ayda Taddayon Tabrizi<sup>3</sup>, Pedram Hashemian<sup>4</sup>, Sajede Alijannejad<sup>5</sup>, Hossein Ali Rahdar<sup>1</sup>, Gordon A Ferns<sup>6</sup>, Seyed Mahdi Hassanian<sup>7</sup>, Soodabeh Shahidsales<sup>8#</sup>, Amir Avan<sup>2,7#</sup>

1. Iranshahr University of Medical Sciences, Iranshahr, Iran.
2. Student Research Committee, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.
3. Master of cellular and Molecular Biology, Genetic Orientation, Faculty of Basic Sciences, Islamic Azad university of Mashhad, Mashhad, Iran.
4. Department of Pathology, Jahad Daneshgahi Institute, Mashhad Branch, Mashhad, Iran.
5. Department of Biology, Faculty of Sciences, Mashhad Branch, Islamic Azad University, Mashhad, Iran.
6. Brighton & Sussex Medical School, Division of Medical Education, Falmer, Brighton, Sussex BN1 9PH, UK.
7. Metabolic Syndrome Research Center, Mashhad University of Medical Sciences, Mashhad, Iran.
8. Cancer Research Center, Mashhad University of Medical Sciences, Mashhad, Iran.

**Running title:** Wnt/  $\beta$ -catenin signaling regulatory microRNAs in breast cancer

Authors have no conflicts of interest.

## **# Corresponding Authors:**

Amir Avan and Soodabeh Shahidsales. Metabolic Syndrome Research Center, Mashhad University of Medical Sciences, Mashhad, Iran. [avana@mums.ac.ir](mailto:avana@mums.ac.ir)  
Shahidsaless@mums.ac.ir

**Funding:** This research was partly supported by Mashhad University of Medical Sciences, (Amir Avan).

**Disclosures:** The authors have no conflicts of interest to declare.

## **Abstract**

Breast cancer is the most commonly diagnosed malignancy in women worldwide. Recently, uncontrolled expression of microRNAs was detected in several human disorders like cardiovascular, neurological, intestinal and autoimmunity diseases. MicroRNAs (miRNAs) are now investigated as novel prognostic and diagnostic biomarkers for several solid tumors like breast, lung, and gastrointestinal cancers. Current data suggest that miRNAs are implicated in various oncogenic processes implicated in breast cancer carcinogenesis through modulating canonical Wnt pathway. Aberrant activation of Wnt/ $\beta$ -catenin signaling was shown to be significantly associated with tumor progression and poor prognosis in patients with breast cancer. This review presents recent findings on the molecular mechanism of microRNAs in regulation of Wnt/ $\beta$ -catenin signaling involved in tumorigenesis of breast cancer.

**Keywords:** Wnt/ $\beta$ -catenin signaling; breast cancer; MicroRNA

## 1. Introduction

Breast cancer ranks as the most prevalent malignancy around the world and the second reason of cancer related death in female population [1]. Despite great improvements in screening, diagnosis and therapeutic strategies, the prognosis and survival outcomes in patients with breast cancer are still unsatisfying [2]. Therefore, it is clinically desirable to identify novel prognostic markers in breast cancer. Recent studies revealed that breast cancer as a heterogeneous disease can be divided into several molecular subtypes with different pathological features and clinical responses [3, 4]. The breast cancer tumorigenesis was recognized as a multistep process which is modulated by various oncogenic cell signaling cascades including PI3K/AKT and Wnt/b-catenin pathways [5-9]. For instance, upregulation of b-catenin and its downstream targets were shown to be significantly related to malignant features and poor clinical outcomes in patients with breast cancer [10, 11]. Recent molecular investigations to identify novel mechanisms involved in breast cancer tumorigenesis indicate that some miRNAs were upregulated along cancer development leading to aberrant activation of various oncogenic signaling pathways.

MicroRNAs (miRNAs) are endogenous, small non-coding RNAs, that are involved in regulating gene expression by pairing with 3'-untranslated regions in target mRNAs. Aberrant expression of multiple miRNAs is considered to have critical roles in controlling various biological events like cell differentiation, metabolism, proliferation, tumorigenesis and drug resistance [12-14]. Regarding their target genes, miRNAs can act as tumor-suppressors or oncogenes in modulating intracellular signalings events. It is well established that the over-expressed miRNAs have prominent effects on tumor cell growth and progression by activating multiple oncogenic signalings like Wnt/ $\beta$ -catenin pathway [15, 16]. The oncogenic activity of these miRNAs is mediated by targeting several tumor suppressors including GSK-3  $\beta$ , APC and PTEN resulted in tumor growth and metastasis in breast cancer patients. Conversely, the tumor suppressor

miRNAs were shown to have anti-cancer role by inhibiting tumor cell proliferation through suppressing  $\beta$ -catenin or Wnt receptors like frizzled proteins.

The regulatory function of miRNAs on the Wnt/ $\beta$ -catenin signaling, suggesting a promising role of miRNAs in breast cancer therapy. Here we reviewed recent findings on the regulatory effects of miRNAs on Wnt/ $\beta$ -catenin signaling involved in tumorigenesis of breast cancer.

## **2. Wnt/ $\beta$ -catenin signaling pathway**

The Wnt/  $\beta$ -catenin signaling plays essential roles in cancer cell growth, survival and metastasis [17, 18]. Generally, the intracellular Wnt signaling subdivided into  $\beta$ -catenin dependent (canonical) and  $\beta$ -catenin in-dependent (non-canonical) routes. The canonical Wnt/ $\beta$ -catenin axis was found to be associated with several oncogenic events including tumor cell proliferation, migration, EMT and invasion [18, 19]. In the condition that canonical Wnt signaling is inactivated, the cytoplasmic  $\beta$ -catenin proteins were phosphorylated by tumor suppressors glycogen synthase kinase 3  $\beta$  (GSK-3  $\beta$ ) and casein kinase 1 (CK1) leading to ubiquitination and proteolysis by proteasomes resulted in inactivation of  $\beta$ -catenin signaling [20]. However, the canonical Wnt signaling was shown to be activated through binding of Wnt ligand to its specific receptors including LRP-5/6 (LDL receptor-related protein 5 or 6) and Frizzled proteins. This in turn induces next components of Wnt pathway comprising dishevelled resulted in deactivation of GSK-3  $\beta$  and dissociation of  $\beta$ -catenin proteins from destruction complex formed by GSK-3  $\beta$ , CK1, Axin, and APC (adenomatous polyposis coli). Therefore,  $\beta$ -catenin proteins accumulated in the cytosol can transferred to the nucleus and regulate multiple downstream genes like c-Myc and cyclin D1[21, 22].

Cyclins, which are considered as key promoters of cellular proliferation, have a prominent role in cell cycle progression through inducing cyclin dependent kinase enzymes [10, 23]. Nowadays, it is well-known that any defect in regulation of  $\beta$ -catenin target genes may causes un-controlled cell cycle progression and development of several types of human tumors like

breast, lymphoma, melanoma, colorectal, and prostate cancers [24-27]. Moreover, prolonged activity of Wnt/ $\beta$ -catenin signaling was shown to be associated with clinical poor outcomes and resistance to chemotherapy [28, 29].

### **3. Regulatory role of oncogenic miRNAs in Wnt/ $\beta$ -catenin signaling in breast cancer cells**

It is widely accepted that over-activation of Wnt/ $\beta$ -catenin signaling is implicated in tumorigenesis and progression of multiple human cancers [18]. Growing body of evidence indicate that downregulation of tumor suppressors involved in regulation of Wnt/ $\beta$ -catenin pathway have critical roles in breast cancer tumorigenesis. As shown in figure 1, there are various studies demonstrated that some miRNAs can promote Wnt signaling through suppressing negative regulators such as GSK-3  $\beta$  and APC [30]. Recently, miR-1229 was detected to be upregulated in breast tumor patients. Aberrant expression of miR-1229 induces Wnt/ $\beta$ -catenin signaling through targeting GSK-3  $\beta$ , APC and ICAT proteins [30]. ICAT also known as inhibitor of  $\beta$ -catenin and TCF (T cell factor), inhibits the expression of downstream targets of Wnt signaling by preventing interaction of  $\beta$ -catenin and TCF. The overexpression of miR-1301 was found to be associated with aggressive features and poor clinical outcomes in patients with breast cancer. It has been demonstrated that the miR-1301 can downregulate ICAT expression by directly binding to 3'-UTR of its mRNA [31]. These data lead to considering miR-1301 as a novel prognostic biomarker for breast tumor. MiR-125b is another Wnt/  $\beta$ -catenin signaling pathway regulatory miRNA, which is overexpressed in breast cancer [32]. It has been shown that the 3'-UTR of APC mRNA have targeted by miR-125b providing the mechanism by which miR-125b can induce Wnt/  $\beta$ -catenin signaling by targeting APC. Downregulation of miR-125b was found to be correlated with inhibition of Wnt/  $\beta$ -catenin signaling resulted in decreased tumor development and dissemination [32]. Thus, repression of miR-125b or overexpression of APC may be considered as potential therapeutic targets for management of breast tumors.

The expression of GSK-3  $\beta$  protein was also found to be inhibited by miR-3646. Moreover, elevated levels of miR-3646 in breast cancer patients was shown to be associated with both docetaxel resistance and upregulation of  $\beta$ -catenin [33]. To further investigate the mechanism of miRNAs in drug resistance in breast cancer, Shen et al. demonstrated that miR-29a was significantly upregulated in Adriamycin-resistant MCF-7 cells [34]. Ectopic expression of miR-29a was observed to be related to downregulation of GSK-3  $\beta$  and PTEN tumor suppressors in MCF-7 breast cancer cells. Therefore, blockade of PTEN/GSK-3  $\beta$  signaling may result in Adriamycin resistance in breast tumors. Taken together, these results indicate that miR-3646 and miR-29a can induce drug resistance in breast tumor cells by promoting Wnt/ $\beta$ -catenin signaling through suppression of GSK-3  $\beta$ .

Similarly, Ma et al. indicated that miR-301 induces Wnt signaling by targeting PTEN (phosphatase and tensin homolog) in breast cancer cells [35]. PTEN was identified as a master regulator of several oncogenic pathways including PI3K/AKT and Wnt signaling which is downregulated in various human cancers [9, 36, 37].

Additionally, previous findings demonstrated that certain miRNAs can induce canonical Wnt pathway through targeting Wnt antagonists like Wnt inhibitory factor-1 (WIF1). WIF1 was shown to be repressed via miR-374a which is upregulated in patients with metastatic breast cancer [38]. The overexpression of miR-374a was reported to be strongly associated with tumor growth and reduced survival in breast cancer patients with distant metastases. To further elucidate the metastatic function of miR-374a, it has been found that miR-374a downregulates the expression of Wnt5a in breast cancer cells [38]. Wnt5a as one of the main regulators of non-canonical pathway, activates CK1 leading to phosphorylation of  $\beta$ -catenin and consequently inhibition of Wnt pathway [39, 40]. These data provide evidence that the oncogenic function of miR-374a in inducing cancer cell metastasis can be mediated by suppressing Wnt/ $\beta$ -catenin signaling inhibitors. To further explore whether oncogenic miRNAs can inhibit regulatory proteins of Wnt/ $\beta$ -catenin signaling, Ren et al. reported that miR-638 is upregulated in breast cancer and

inhibits DACT3 as one of the main modulators of Wnt/  $\beta$ -catenin pathway [41]. DACT3, a member of DACT family, was identified as a potent tumor suppressor which inhibits tumor cell proliferation by regulating  $\beta$ -catenin expression in various human malignancies [42]. Additionally, it has been reported that upregulation of miR-638 was directly correlated with malignant features of breast cancer while regulates cellular autophagy [41].

In another investigation, Chen et al. demonstrated that aberrant expression of miR-373 has prominent role in promoting EMT by inducing mesenchymal markers like  $\beta$ -catenin in breast cancer cells [43]. These results suggest that miR-373 can promote epithelial-to-mesenchymal transition (EMT) and metastasis in breast tumors by upregulating Wnt/  $\beta$ -catenin signaling pathway.

#### **4. Regulatory role of tumor suppressor miRNAs in Wnt/ $\beta$ -catenin signaling in breast cancer cells**

$\beta$ -catenin as a potent oncogene is upregulated in multiple human cancers.  $\beta$ -catenin regulates expression of multiple proteins involved in cell proliferation like cyclin D1 which is found to be upregulated in nearly 50% of patients with breast cancer [10]. Aberrant activity of  $\beta$ -catenin was directly related to malignant manifestations and poor prognosis in breast cancer [10, 44].  $\beta$ -catenin was identified as target gene for some regulatory miRNAs including miR-26b, miR-135, miR-214, miR-216a, and miR-340 that are downregulated in breast tumor cells [45-49]. More investigations on the mechanism of miR-135 was also demonstrated that this tumor suppressor miRNA can inhibit tumor cell proliferation and EMT by regulating GSK3-  $\beta$  and matrix metalloproteinase (MMP) enzymes respectively [46]. Consistently, Xiong et al. reported that miR-30a-5p represses  $\beta$ -catenin downstream targets like cyclin B1, cyclin D1 and c-Myc in breast tumor cells [50]. It has been also reported that expression of miR-30a-5p is inversely related to the expression of ubiquitin protein ligase E3C (UBE3C) in breast cancer cell lines. UBE3C was



shown to be upregulated in breast cancer leads to overexpression of c-Myc and cyclin D1 [50, 51].

Furthermore, miR-138 and miR-449b-5p were identified as tumor suppressor miRNAs which inhibit tumor progression and metastasis by targeting CREPT (Cell Cycle-Related and Expression-elevated Protein in Tumor) [52, 53]. The overexpression of CREPT was detected in breast tumor cells and tissues. CREPT as an oncogene protein, has a critical role in cellular proliferation through modulating cell cycle-related genes like cyclin D1 [54, 55]. MiR-384 is another tumor suppressor microRNA which is downregulated in breast cancer patients [56]. Upregulation of miR-384 inhibits ACVR1 (Activin A receptor type 1) resulting in repression of Wnt/ $\beta$ -catenin axis in breast tumors. Previous data indicate that ACVR1 can induce tumor cell metastasis by promoting Wnt/ $\beta$ -catenin pathway in several human malignancies like breast cancer [57, 58]. Consistent with these findings, it has been shown that miR-30 inhibits cancer cell invasion and metastasis while induces cell apoptosis by repressing CTHRC1 (Collagen triple helix repeat containing-1) in breast tumors [59]. CTHRC1 is a secretory protein involved in stabilizing binding of Wnt ligand to their Frizzled receptors leading to activation of Wnt signaling pathway. Further studies on the oncogenic function of CTHRC1 indicate that elevated levels of CTHRC1 induces GSK3  $\beta$  and  $\beta$ -catenin activities which are associated with aggressive clinicopathological manifestations and worse prognosis in several malignant tumors including gastrointestinal, liver, breast and non-small cell lung cancers [60-62]. MiR-140-5p is another Wnt inhibitory miRNA which is down-regulated in breast cancer stem cells. Ectopic expression of miR-140-5p represents anti-tumor activity mediated by targeting Wnt1 [63].

To further investigate the inhibitory function of regulatory miRNAs on Wnt/  $\beta$ -catenin axis, it has been reported that some miRNAs can limit tumor progression by targeting Wnt receptors like Frizzled proteins. For instance, Liu et al. reported that miR-224 represents anti-cancer properties through targeting Frizzled 5 receptor proteins resulting in inhibition of Wnt signaling in breast cancer [64]. In line with this, Leu et al. demonstrated that miR-1 inhibits Frizzled 7 and

tankyrase 2 proteins resulted in reduced cellular proliferation and migration in breast tumors [65]. Similarly, Jiang et al illustrated that miR-100 which is repressed in breast cancer patients can inhibit Wnt/  $\beta$ -catenin pathway by directly targeting Frizzled 8 receptor proteins [66]. In another study, the effective role of regulatory miRNAs on secreted Frizzled-related proteins (SFRPs) was investigated in breast tumor cells. Their results indicate that miR-27a can restrict tumor cell proliferation, migration and invasion by targeting SFRPs [67]. These results support the idea that Frizzled receptors and their related proteins may be considered as promising targets for breast cancer therapy.

To further identify the oncogenic targets regulated by miRNAs, it has been shown that cyclin dependent kinase enzyme 8 (CDK8) has been downregulated by certain miRNAs including miR-107 and miR-26b [45, 68]. The CDK8 enzyme has critical role in promoting cell cycle progression and cellular proliferation resulted in tumor growth and development.

## **5. Conclusion**

Recent data indicate that multiple miRNAs are differentially expressed in breast cancer, and have critical roles in tumor cell proliferation, invasion, and metastasis (Table 1 and 2).

As shown in figure 2, the overexpression of various miRNAs including miR-1229, miR-125b, miR-3646, miR-29a, and miR-301 might contribute to aberrant activation of the Wnt/  $\beta$ -catenin signaling. The oncogenic activity of these miRNAs is mediated by targeting several tumor suppressors including GSK-3  $\beta$ , APC and PTEN. Consistently, the Wnt signaling downstream target genes were upregulated resulted in tumor growth, metastasis and drug resistance in breast cancer patients. Conversely, the tumor suppressor miRNAs including miR-26b, miR-135, miR-214, miR-216a, and miR-340 were found to have downregulated in breast cancer patients. These miRNAs were shown to have anti-tumor activity by inhibiting cancer cell proliferation through suppressing  $\beta$ -catenin. These data in combination with recent findings demonstrate that

uncontrolled activity of oncogenic Wnt/  $\beta$ -catenin signaling is associated to poor prognosis and aggressive behavior in patients with breast cancer.

In this review, we briefly discussed the previous data about the function of Wnt axis regulatory miRNAs involved in carcinogenesis of breast cancer. At present, there is a growing trend on targeting oncogenic miRNAs by certain inhibitors for anti-cancer approaches. Considering the oncogenic or anti-tumor function of miRNAs and their target genes, it has been proposed that miRNAs may be considered as new diagnostic and prognostic biomarkers for breast cancer. Therefore, regulating the oncogenic miRNAs by specific inhibitors or upregulation of anti-tumor miRNAs may be considered as novel miRNA-based therapeutic strategies for management of breast cancer.

## References

1. Zaidi, Z. and H.A. Dib, *The worldwide female breast cancer incidence and survival*, 2018. 2019, AACR.
2. McKinney, S.M., et al., *International evaluation of an AI system for breast cancer screening*. *Nature*, 2020. **577**(7788): p. 89-94.
3. Søkilde, R., et al., *Refinement of breast cancer molecular classification by miRNA expression profiles*. *BMC genomics*, 2019. **20**(1): p. 503.
4. Zhang, L., et al., *Comparison of breast cancer risk factors among molecular subtypes: A case-only study*. *Cancer medicine*, 2019. **8**(4): p. 1882-1892.
5. Owusu-Brackett, N., M. Shariati, and F. Meric-Bernstam, *Role of PI3K/AKT/mTOR in Cancer Signaling*, in *Predictive Biomarkers in Oncology*. 2019, Springer. p. 263-270.
6. Sharma, V., et al. *Recent nanotechnological interventions targeting PI3K/Akt/mTOR pathway: A focus on breast cancer*. in *Seminars in cancer biology*. 2019. Elsevier.
7. Khan, M.A., et al., *PI3K/AKT/mTOR pathway inhibitors in triple-negative breast cancer: a review on drug discovery and future challenges*. *Drug discovery today*, 2019.
8. Krishnamurthy, N. and R. Kurzrock, *Targeting the Wnt/beta-catenin pathway in cancer: Update on effectors and inhibitors*. *Cancer treatment reviews*, 2018. **62**: p. 50-60.
9. Rahmani, F., et al., *Role of regulatory miRNAs of the PI3K/AKT signaling pathway in the pathogenesis of breast cancer*. *Gene*, 2020: p. 144459.
10. Lin, S.-Y., et al.,  *$\beta$ -catenin, a novel prognostic marker for breast cancer: its roles in cyclin D1 expression and cancer progression*. *Proceedings of the National Academy of Sciences*, 2000. **97**(8): p. 4262-4266.
11. Pang, H., et al., *Prognostic values of osteopontin-c, E-cadherin and  $\beta$ -catenin in breast cancer*. *Cancer epidemiology*, 2013. **37**(6): p. 985-992.
12. Hwang, H. and J. Mendell, *MicroRNAs in cell proliferation, cell death, and tumorigenesis*. *British journal of cancer*, 2006. **94**(6): p. 776-780.
13. Aravindan, N., et al., *MicroRNAs in neuroblastoma tumorigenesis, therapy resistance, and disease evolution*. *Cancer drug resistance (Alhambra, Calif.)*, 2019. **2**: p. 1086.
14. Soleimani, A., et al., *The potential role of regulatory microRNAs of RAS/MAPK signaling pathway in the pathogenesis of colorectal cancer*. *Journal of cellular biochemistry*, 2019. **120**(12): p. 19245-19253.
15. Slattery, M.L., et al., *Expression of Wnt-signaling pathway genes and their associations with miRNAs in colorectal cancer*. *Oncotarget*, 2018. **9**(5): p. 6075.
16. LIN, H., et al., *Regulatory mechanism of microRNA and Wnt/ $\beta$ -catenin signaling pathway in tumorigenesis and metastasis*. *SCIENTIA SINICA Vitae*, 2018. **48**(1): p. 5-14.
17. Li, G., et al., *Frizzled7 promotes epithelial-to-mesenchymal transition and stemness via activating canonical Wnt/ $\beta$ -catenin pathway in gastric cancer*. *International journal of biological sciences*, 2018. **14**(3): p. 280.
18. Rahmani, F., et al., *Role of Wnt/ $\beta$ -catenin signaling regulatory microRNAs in the pathogenesis of colorectal cancer*. *Journal of cellular physiology*, 2018. **233**(2): p. 811-817.
19. Rahmani, F., et al., *PNU-74654 enhances the antiproliferative effects of 5-FU in breast cancer and antagonizes thrombin-induced cell growth via the Wnt pathway*. *Journal of cellular physiology*, 2019. **234**(8): p. 14123-14132.
20. Song, J.L., et al., *microRNA regulation of Wnt signaling pathways in development and disease*. *Cellular signalling*, 2015. **27**(7): p. 1380-1391.

21. Gustafson, C.T., et al., *Molecular strategies for modulating Wnt signaling*. Front Biosci (Landmark Ed), 2017. **22**: p. 137-156.
22. Anastas, J.N. and R.T. Moon, *WNT signalling pathways as therapeutic targets in cancer*. Nature Reviews Cancer, 2013. **13**(1): p. 11-26.
23. Huo, D., et al., *Omethoate induces pharyngeal cancer cell proliferation and G1/S cell cycle progression by activation of Akt/GSK-3 $\beta$ /cyclin D1 signaling pathway*. Toxicology, 2019. **427**: p. 152298.
24. Zhao, Y., et al., *Cyclin D1 overexpression is associated with poor clinicopathological outcome and survival in oral squamous cell carcinoma in Asian populations: insights from a meta-analysis*. PloS one, 2014. **9**(3).
25. Gwak, H., et al., *Metformin induces degradation of cyclin D1 via AMPK/GSK3 $\beta$  axis in ovarian cancer*. Molecular carcinogenesis, 2017. **56**(2): p. 349-358.
26. Casimiro, M.C., et al., *Overview of cyclins D1 function in cancer and the CDK inhibitor landscape: past and present*. Expert opinion on investigational drugs, 2014. **23**(3): p. 295-304.
27. Prasad, A., et al., *Styryl sulfonyl compounds inhibit translation of cyclin D1 in mantle cell lymphoma cells*. Oncogene, 2009. **28**(12): p. 1518-1528.
28. Zhang, H., et al., *Interference of Frizzled 1 (FZD1) reverses multidrug resistance in breast cancer cells through the Wnt/ $\beta$ -catenin pathway*. Cancer letters, 2012. **323**(1): p. 106-113.
29. Nagaraj, A.B., et al., *Critical role of Wnt/ $\beta$ -catenin signaling in driving epithelial ovarian cancer platinum resistance*. Oncotarget, 2015. **6**(27): p. 23720.
30. Tan, Z., et al., *MicroRNA-1229 overexpression promotes cell proliferation and tumorigenicity and activates Wnt/ $\beta$ -catenin signaling in breast cancer*. Oncotarget, 2016. **7**(17): p. 24076.
31. Lin, W.-h., et al., *MicroRNA-1301 induces cell proliferation by downregulating ICAT expression in breast cancer*. Biomedicine & Pharmacotherapy, 2016. **83**: p. 177-185.
32. Nie, J., et al., *Mir-125b regulates the proliferation and metastasis of triple negative breast cancer cells via the Wnt/ $\beta$ -catenin pathway and EMT*. Bioscience, Biotechnology, and Biochemistry, 2019. **83**(6): p. 1062-1071.
33. Zhang, X., et al., *MicroRNA-3646 contributes to docetaxel resistance in human breast cancer cells by GSK-3 $\beta$ / $\beta$ -catenin signaling pathway*. PloS one, 2016. **11**(4).
34. Shen, H., et al., *MicroRNA-29a contributes to drug-resistance of breast cancer cells to adriamycin through PTEN/AKT/GSK3 $\beta$  signaling pathway*. Gene, 2016. **593**(1): p. 84-90.
35. Ma, F., et al., *Upregulated microRNA-301a in breast cancer promotes tumor metastasis by targeting PTEN and activating Wnt/ $\beta$ -catenin signaling*. Gene, 2014. **535**(2): p. 191-197.
36. Liu, Y.-Z., et al., *The PTEN/PI3K/Akt and Wnt/ $\beta$ -catenin signaling pathways are involved in the inhibitory effect of resveratrol on human colon cancer cell proliferation*. International journal of oncology, 2014. **45**(1): p. 104-112.
37. Perry, J.M., et al., *Cooperation between both Wnt/ $\beta$ -catenin and PTEN/PI3K/Akt signaling promotes primitive hematopoietic stem cell self-renewal and expansion*. Genes & development, 2011. **25**(18): p. 1928-1942.
38. Cai, J., et al., *MicroRNA-374a activates Wnt/ $\beta$ -catenin signaling to promote breast cancer metastasis*. The Journal of clinical investigation, 2013. **123**(2).
39. Foldynová-Trantírková, S., et al., *Breast cancer-specific mutations in CK1 $\epsilon$  inhibit Wnt/ $\beta$ -catenin and activate the Wnt/Rac1/JNK and NFAT pathways to decrease cell adhesion and promote cell migration*. Breast Cancer Research, 2010. **12**(3): p. R30.
40. Medrek, C., et al., *Wnt-5a-CK1 $\alpha$  signaling promotes  $\beta$ -catenin/E-cadherin complex formation and intercellular adhesion in human breast epithelial cells*. Journal of Biological Chemistry, 2009. **284**(16): p. 10968-10979.

41. Ren, Y., et al., *MiRNA-638 promotes autophagy and malignant phenotypes of cancer cells via directly suppressing DACT3*. Cancer letters, 2017. **390**: p. 126-136.
42. Jiang, X., et al., *DACT3 is an epigenetic regulator of Wnt/ $\beta$ -catenin signaling in colorectal cancer and is a therapeutic target of histone modifications*. Cancer cell, 2008. **13**(6): p. 529-541.
43. De Chen, B.-L.D., et al., *MiR-373 drives the epithelial-to-mesenchymal transition and metastasis via the miR-373-TXNIP-HIF1 $\alpha$ -TWIST signaling axis in breast cancer*. Oncotarget, 2015. **6**(32): p. 32701.
44. López-Knowles, E., et al., *Cytoplasmic localization of  $\beta$ -catenin is a marker of poor outcome in breast cancer patients*. Cancer Epidemiology and Prevention Biomarkers, 2010. **19**(1): p. 301-309.
45. Li, J., et al., *MiRNA-26b inhibits cellular proliferation by targeting CDK8 in breast cancer*. International journal of clinical and experimental medicine, 2014. **7**(3): p. 558.
46. Jiang, D., et al., *miR-135 regulated breast cancer proliferation and epithelial-mesenchymal transition acts by the Wnt/ $\beta$ -catenin signaling pathway*. International journal of molecular medicine, 2019. **43**(4): p. 1623-1634.
47. Yi, S., L. Li, and W. Tu, *MiR-214 negatively regulates proliferation and WNT/ $\beta$ -catenin signaling in breast cancer*. Eur Rev Med Pharmacol Sci, 2016. **20**(24): p. 5148-5154.
48. Xie, Q., et al., *MicroRNA-216a suppresses the proliferation and migration of human breast cancer cells via the Wnt/ $\beta$ -catenin signaling pathway*. Oncology reports, 2019. **41**(5): p. 2647-2656.
49. Mohammadi-Yeganeh, S., et al., *MicroRNA-340 inhibits the migration, invasion, and metastasis of breast cancer cells by targeting Wnt pathway*. Tumor Biology, 2016. **37**(7): p. 8993-9000.
50. Xiong, J., et al., *MiR-30a-5p/UBE3C axis regulates breast cancer cell proliferation and migration*. Biochemical and biophysical research communications, 2019. **516**(3): p. 1013-1018.
51. Wen, J.L., et al., *UBE3C promotes growth and metastasis of renal cell carcinoma via activating Wnt/ $\beta$ -catenin pathway*. PloS one, 2015. **10**(2).
52. Liang, Z., et al., *CREPT regulated by miR-138 promotes breast cancer progression*. Biochemical and biophysical research communications, 2017. **493**(1): p. 263-269.
53. Jiang, J., et al., *MicroRNA-449b-5p suppresses the growth and invasion of breast cancer cells via inhibiting CREPT-mediated Wnt/ $\beta$ -catenin signaling*. Chemico-biological interactions, 2019. **302**: p. 74-82.
54. Ma, J., et al., *Knocking-down of CREPT prohibits the progression of oral squamous cell carcinoma and suppresses cyclin D1 and c-Myc expression*. PloS one, 2017. **12**(4).
55. Liu, T., et al., *Inhibiting CREPT reduces the proliferation and migration of non-small cell lung cancer cells by down-regulating cell cycle related protein*. American journal of translational research, 2016. **8**(5): p. 2097.
56. Wang, Y., Z. Zhang, and J. Wang, *MicroRNA-384 inhibits the progression of breast cancer by targeting ACVR1*. Oncology reports, 2018. **39**(6): p. 2563-2574.
57. Li, L., et al., *Regulatory MiR-148a-ACVR1/BMP circuit defines a cancer stem cell-like aggressive subtype of hepatocellular carcinoma*. Hepatology, 2015. **61**(2): p. 574-584.
58. Wang, K., et al., *DNALK2 inhibits the proliferation and invasiveness of breast cancer MDA-MB-231 cells through the Smad-dependent pathway*. Oncology reports, 2017. **37**(2): p. 879-886.
59. Lai, Y.-h., et al., *Collagen triple helix repeat containing-1 negatively regulated by microRNA-30c promotes cell proliferation and metastasis and indicates poor prognosis in breast cancer*. Journal of Experimental & Clinical Cancer Research, 2017. **36**(1): p. 92.
60. Tang, L., et al., *Aberrant expression of collagen triple helix repeat containing 1 in human solid cancers*. Clinical cancer research, 2006. **12**(12): p. 3716-3722.
61. Ke, Z., et al., *Overexpression of collagen triple helix repeat containing 1 (CTHRC1) is associated with tumour aggressiveness and poor prognosis in human non-small cell lung cancer*. Oncotarget, 2014. **5**(19): p. 9410.

62. Hou, M., et al., *High expression of CTHRC1 promotes EMT of epithelial ovarian cancer (EOC) and is associated with poor prognosis*. Oncotarget, 2015. **6**(34): p. 35813.
63. Wu, D., et al., *miR-140-5p inhibits the proliferation and enhances the efficacy of doxorubicin to breast cancer stem cells by targeting Wnt1*. Cancer gene therapy, 2019. **26**(3): p. 74-82.
64. Liu, F., et al., *MicroRNA-224 inhibits proliferation and migration of breast cancer cells by down-regulating fizzled 5 expression*. Oncotarget, 2016. **7**(31): p. 49130.
65. Liu, T., et al., *MicroRNA-1 down-regulates proliferation and migration of breast cancer stem cells by inhibiting the Wnt/ $\beta$ -catenin pathway*. Oncotarget, 2015. **6**(39): p. 41638.
66. Jiang, Q., et al., *MicroRNA-100 suppresses the migration and invasion of breast cancer cells by targeting FZD-8 and inhibiting Wnt/ $\beta$ -catenin signaling pathway*. Tumor Biology, 2016. **37**(4): p. 5001-5011.
67. Kong, L.-Y., et al., *In vivo and in vitro effects of microRNA-27a on proliferation, migration and invasion of breast cancer cells through targeting of SFRP1 gene via Wnt/ $\beta$ -catenin signaling pathway*. Oncotarget, 2017. **8**(9): p. 15507.
68. Li, X.-Y., et al., *MiRNA-107 inhibits proliferation and migration by targeting CDK8 in breast cancer*. International journal of clinical and experimental medicine, 2014. **7**(1): p. 32.

## Figures




Figure 1) Schematic model of the role of Wnt/ $\beta$ -catenin signaling regulatory microRNAs in breast cancer tumorigenesis.






Figure 2) The mechanisms of Wnt/  $\beta$ -catenin signaling regulatory miRNAs in breast cancer carcinogenesis.

Table1. Oncogenic miRNAs regulate the pathogenesis of breast cancer by targeting the certain components of the Wnt/ $\beta$ -catenin signaling pathway.

<b>microRNA</b>	<b>Molecular alteration</b>	<b>Target</b>	<b>Function</b>	<b>Reference</b>
miR-1229	Upregulation	GSK-3 $\beta$ , APC, and ICAT	Cellular proliferation	Tan et al., 2016
miR-1301	Upregulation	ICAT	Cellular proliferation	Lin et al., 2016
miR-125b	Upregulation	APC	Cellular proliferation and metastasis	Nie et al., 2019
miR-3646	Upregulation	GSK-3 $\beta$	Drug resistance	Zhang et al., 2016
miR-29a	Upregulation	GSK-3 $\beta$	Drug resistance	Shen et al., 2016
miR-301	Upregulation	PTEN	Cellular proliferation and metastasis	Ma et al., 2014
miR-374a	Upregulation	WIF1 and Wnt5a	Tumor metastasis	Cai et al., 2013
miR-638	Upregulation	DACT3	Cellular proliferation	Ren et al., 2017
miR-373	Upregulation	$\beta$ -catenin	EMT	De Chen et al., 2015

Table2. Tumor suppressor miRNAs regulate the pathogenesis of breast cancer by targeting the certain components of the Wnt/ $\beta$ -catenin signaling pathway.

<b>microRNA</b>	<b>Molecular alteration</b>	<b>Target</b>	<b>Function</b>	<b>Reference</b>
miR-26b	Downregulation	$\beta$ -catenin	Inhibits cellular proliferation	Li et al., 2014a
miR-135	Downregulation	$\beta$ -catenin	Inhibits cellular proliferation	Jiang et al., 2019a
miR-214	Downregulation	$\beta$ -catenin	Inhibits cellular proliferation	Yi et al., 2016
miR-216a	Downregulation	$\beta$ -catenin	Inhibits cellular proliferation	Xie et al., 2019
miR-340	Downregulation	$\beta$ -catenin	Inhibits cellular proliferation	Mohammadi-Yeganeh et al., 2016
miR-30a-5p	Downregulation	UBE3C	Inhibits cellular proliferation and metastasis	Xiong et al., 2019
miR-138	Downregulation	CREPT	Inhibits tumor growth and metastasis	Liang et al., 2017
miR-449b-5p	Downregulation	CREPT	Inhibits tumor growth and metastasis	Jiang et al., 2019
miR-384	Downregulation	ACVR1	Inhibits tumor metastasis	Wang et al., 2018
miR-30	Downregulation	CTHRC1	Inhibits cell invasion and metastasis	Lai et al., 2017
miR-140-5p	Downregulation	Wnt1	Inhibits cellular proliferation	Wu et al., 2019
miR-224	Downregulation	Frizzled 5	Inhibits tumor progression	Liu et al., 2016
miR-1	Downregulation	Frizzled 7	Inhibits tumor progression	Liu et al., 2015
miR-100	Downregulation	Frizzled 8	Inhibits tumor progression	Jiang et al., 2016
miR-27a	Downregulation	SFRPs	Inhibits cell growth and invasion	Kong et al., 2017
miR-107	Downregulation	CDK8	Inhibits cellular proliferation and metastasis	Li et al., 2014
miR-26b	Downregulation	CDK8	Inhibits cellular proliferation and metastasis	Li et al., 2014